



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTIONS OF EXERCISES.

166

A CIRCLE cuts a parabola, and the centroid of the four points of intersection is found. What is the locus of the centre of the circle if this point be fixed?

[*W. M. Thornton.*]

SOLUTION.

The equations to the curves are

$$y^2 = 2px, \quad (x - u)^2 + (y - v)^2 = r^2.$$

At the points of intersection

$$[(x - u)^2 - r^2 + 2px + v^2]^2 = 8pv^2x,$$

or

$$x^4 + 4(p - u)x^3 + \dots = 0.$$

By hypothesis, $x_1 + x_2 + x_3 + x_4$ is constant; or

$$p - u = c,$$

which is a straight line at right angles to the axis of the parabola.

[*R. H. Graves.*]

167

If $y_1 y_2 = k^2$ (a constant), where (x_1, y_1) and (x_2, y_2) are points on the parabola $y^2 = 4ax$, find the locus of the intersections of the normals at these points.

[*R. H. Graves.*]

SOLUTION.

If (u, v) be the foot of the normal from (x, y) to $y^2 = 2px$

$$v^3 + 2p(p - x)v - 2p^2y = 0.$$

By hypothesis $v_1 v_2 = k^2$. But $v_0 v_1 v_2 = 2p^2 y$. Therefore

$$v_0 = \frac{2p^2 y}{k^2},$$

whence we find for the equation to the required locus

$$4p^4 y^2 = 2pk^4(x - p) + k^6.$$

[*W. M. Thornton, and others.*]

205

A TRIANGLE PQR is inscribed in an ellipse having the two sides PQ and PR passing through the foci, and the line QR produced meets the tangent at P at the point S . The polar of S with respect to a concentric circle through P passes through the centre of curvature of the ellipse at the point P .

[*R. H. Graves.*]

SOLUTION.

Let
$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

be the equation to the ellipse, and let the co-ordinates of the points P and S be (x', y') and (x_1, y_1) .

Then, S is the pole of the normal at P . (See solution to Exercise 139.)

$$\therefore x_1 = \frac{a^4}{(a^2 - b^2)x'} \text{ and } y_1 = -\frac{b^4}{(a^2 - b^2)y'}.$$

The centre of curvature is given by the equations

$$\alpha = \frac{(a^2 - b^2)x'^3}{a^4} \text{ and } \beta = -\frac{(a^2 - b^2)y'^3}{b^4};$$

$$\therefore \alpha x_1 + \beta y_1 = x'^2 + y'^2,$$

and the truth of the proposition follows.

[*R. H. Graves.*]

206

IF from any point four normals be drawn to each of the hyperbolas,

$$x^2 - y^2 = a^2 \quad \text{and} \quad xy = k^2,$$

the centres of mean position of the feet of the two sets of normals are coincident.

[*R. H. Graves.*]

SOLUTION.

The normal from (x', y') to $x^2 - y^2 = a^2$ at (x, y) is

$$xy' + x'y = 2xy;$$

$$\therefore (x^2 - a^2)(4x^2 - 4xx' + x'^2) = x^2y'^2;$$

$$\therefore x_1 + x_2 + x_3 + x_4 = x'.$$

A like result will be found for $xy = k^2$.

214

AN equilateral hyperbola is circumscribed to a triangle. Find the greatest and least of the values of the transverse axis. [R. H. Graves.]

SOLUTION.

Since the curve passes through the orthocentre, the minimum is zero, and the maximum is two-thirds of the altitude. [R. H. Graves.]

217

A CIRCLE is drawn at random within a given circle. What is the probability that the random circle contains the centre of the given circle? [Artemas Martin.]

SOLUTION.

Let A be the centre of the given circle, and P the centre of the random circle; through P draw the radius OB of the given circle; then OP is the distance between their centres. Put r = radius of the given circle, and let $OP = x$. In order that the random circle may contain the centre of the given circle, OP , the distance between their centres must be less than $\frac{1}{2}r$. While the centre of the random circle is at P , its radius can have any value from 0 to $r - x$; but it will not contain the centre of the given circle unless its radius exceeds x ; hence in order to include the centre of the given circle, the radius of the random circle can have any value from x to $r - x$; therefore for an assigned value of x the probability that the random circle contains the centre of the given circle is

$$p_x = \frac{r - 2x}{r - x}.$$

But x may have any value from 0 to $\frac{1}{2}r$, and all these values are equally probable. P will be at the distance x from the centre of the given circle if it is anywhere on the circumference of the circle whose centre is at O and radius x ; therefore the probability required is

$$\begin{aligned} p &= \int_0^{\frac{1}{2}r} p_x \times 2\pi x dx \div \int_0^r 2\pi x dx, \\ &= \frac{2}{r^2} \int_0^{\frac{1}{2}r} x \left(\frac{r - 2x}{r - x} \right) dx, \\ &= \frac{2}{r^2} \int_0^{\frac{1}{2}r} \left(2x + r - \frac{r^2}{r - x} \right) dx, \\ &= 2 \left(\frac{3}{4} - \log_e 2 \right). \end{aligned}$$

[Artemas Martin.]